

Current Approaches to Pediatric Polyposis Syndromes

Aodhnait S. Fahy, BMBCh, PhD¹ Christopher R. Moir, MD¹

¹Division of Pediatric Surgery, Department of Surgery, Mayo Clinic, Rochester, Minnesota

Clin Colon Rectal Surg 2018;31:132–142.

Address for correspondence Christopher R. Moir, MD, Division of Pediatric Surgery, Department of Surgery, Mayo Clinic, 200 First Street SW, Rochester, MN 55905
(e-mail: moir.christopher@mayo.edu).

Abstract

Keywords

- IPAA
- pediatric polyposis
- restorative proctocolectomy

Colorectal adenomatous polyposis syndromes encompass a diverse group of disorders with varying modes of inheritance and penetrance. Children may present with overt disease or within screening programs for families at high risk. We provide an overview of the array of pediatric polyposis syndromes, current screening recommendations, and surgical indications and technical considerations. Optimal disease management for these pediatric patients is still evolving and has implications for screening, surveillance, pediatric surgical management, and transition of care gastroenterologic neoplasia physicians and surgeons.

Polyp syndromes derive from adenomatous, hamartomatous, or hyperplastic polyps. Treatment varies significantly based on the nature and distribution of polyps, disease volume, risk of malignant transformation, and type of gastroenterologic or surgical management.

Adenomatous Polyp Syndromes

Familial Adenomatous Polyposis

Familial adenomatous polyposis (FAP) is the most common of the polyposis syndromes, with an estimated prevalence of 2 to 3 per 100,000 persons across all ethnic groups.¹ It is an autosomal dominantly inherited cancer syndrome characterized by mutations in the gene for adenomatous polyposis coli (APC) and development of hundreds to thousands of colorectal adenomatous polyps.² FAP is defined clinically as ≥ 100 adenomatous colon and rectal polyps. It typically presents with malignancy or symptoms in patients younger than 40 years. Left untreated, the FAP penetrance for colorectal cancer (CRC) is close to 100% by the age of 50 years.³

The APC gene spans 108 kb of genomic DNA on chromosome 5.³ It encodes a tumor suppressor protein that functions by downregulating the β -catenin oncoprotein. In the absence of the APC protein, β -catenin accumulates in the nucleus and leads to the involvement of upregulation of genes in cell cycle entry and dysplasia. More than 400 mutations of the APC gene correlate with the clinical phe-

notype and the severity of manifestations. Classic FAP is associated with other benign and premalignant lesions throughout the gastrointestinal tract, including fundic gland gastric polyps, gastric adenomatous polyps, and small-bowel polyps, as well as periampullary carcinoma.⁴

Attenuated Familial Adenomatous Polyposis Syndromes

Several variants of FAP have different genetic mutations associated with various ages of onset and various CRC risks.

Patients with attenuated AFAP usually have fewer polyps that occur at a later age of onset. In addition, rectal sparing and right-sided adenomas often develop. These patients do not have extracolonic mutations and have a reduced risk of CRC. AFAP arises from mutations in the APC gene at either the proximal or distal end or in locations of exon 9.⁵

MUTYH-associated polyposis is the second most common attenuated FAP-related syndrome and involves the MUTYH gene—which is active in the repair of DNA damaged by oxidative stress—thereby resulting in transversions in multiple genes, including APC and KRAS.⁶ Biallelic germline mutations in MUTYH are known to cause a less severe form of adenomatous polyposis, similar to attenuated FAP, in which the patient typically has less than 100 adenomatous polyps. MUTYH-associated polyposis is inherited in an autosomal recessive way, and MUTYH mutations have been identified in 7.5 to 12.5% of patients who have more than 100 adenomas without an APC mutation.

Polymerase proofreading–associated polyposis is a recently described syndrome where the phenotype includes a reduced number of polyps (relative to *APC* or *MUTYH*) but with high penetrance, high dysplastic risk, and early progression to CRC.⁷ Germline mutations in enzymes encoding the catalytic and proofreading activities of the leading-strand DNA polymerase epsilon (*POLE*) and the lagging-strand polymerase delta (*POLD*) are responsible.⁸ Polymerase proofreading–associated polyposis has been reported in up to 7% of families with colorectal adenomas and carcinomas without *APC* or *MUTYH* mutations.⁹

In 30 to 50% of patients with adenomatous polyposis, no germline mutation can be identified in the *APC*, *MUTYH*, or *POLE/POLD1* genes. An *APC* mutational mosaicism may be detected in 11 to 20% of patients. An additional 30 low-susceptibility variants have been identified that increase risk of CRC.⁹ Whether a patient carries a genetic susceptibility for cancer is an important determinant of when and how to screen the patient, as well as an indication for their likely response to chemotherapy, and their expected prognosis. Patients with mutational mosaicism or low-susceptibility variants typically received the diagnosis of malignancy at an older age than those who were clear gene carriers, though many of these patients with low susceptibility variants or mutational mosaicism had a strong family history of CRC, conforming to Amsterdam II criteria,¹⁰ and as such were still candidates for prophylactic surgical management of their disease process.

Hamartomatous Polyp Syndromes

Hamartomatous polyps are pedunculated, cherry-red polypoid lesions. Hamartomatous polyp syndromes that increase rates of CRC include Peutz-Jeghers syndrome (PJS), juvenile polyposis syndrome (JPS), and phosphatase tension homolog (*PTEN*) hamartoma syndrome.

PJS is an autosomal dominant syndrome. Diagnostic criteria include the presence of ≥ 2 hamartomatous polyps anywhere in the gastrointestinal tract, one confirmed hamartomatous polyp in a patient with a family history of PJS, and typical perioral pigmentation.¹¹ The mutation occurs in a gene that codes for a serine or threonine kinase (*LKB1* or *STK11*), which acts as a tumor suppressor. Varying mutations lead to the phenotypic variability of PJS. The lifetime risk of CRC in PJS has been estimated as 39%, and PJS patients are also vulnerable to pancreatic cancers, breast tumors, and cervical and testicular tumors.^{12,13} Management of PJS centers on resection of large polyps and surveillance for malignant tumors. Patients should undergo upper and lower endoscopy biennially after the age of 30 years, and endoscopic resection is recommended for polyps greater than 1.5 cm. Family members also should undergo regular screening.¹⁴

Juvenile polyposis syndrome is a rare syndrome inherited with incomplete penetrance in 20 to 50% patients with hamartomatous polyps. JPS diagnostic criteria include more than five juvenile polyps in the colon or the rectum, multiple juvenile polyps throughout the entire gastrointestinal tract, and one or more polyps in combination with a family history of juvenile polyposis.¹⁵ Germline mutations have been described in *SMAD4* and *BMPR1* genes in some, but not all, cases of JPS. The most affected sites of polyps in JPS

are colon and rectum (98%), stomach (14%), jejunum and ileum (7%), and duodenum (2%).¹⁵ The estimated CRC risk is 17 to 22% by the age of 35 years, though gastric, duodenal, and pancreatic cancers are also common.¹⁶ Depending on location, juvenile polyps should be resected endoscopically or surgically. A prophylactic colectomy has been advocated for patients who have polyps with early adenomatous features, dysplasia, or a family history of CRC.¹⁷ Surveillance of the pouch and ileal mucosa should be ongoing.¹⁸

PTEN hamartomatous tumor syndromes (PHTSs) include Cowden's syndrome and Bannayan-Riley-Ruvalcaba syndrome. Both of these syndromes are characterized by the autosomal dominant inheritance of mutations in the *PTEN* (phosphatase and tensin homolog) gene, which encodes a tumor suppressor. *PTEN* mutations are found in 25% of patients with PHTS; other patients have *SDH* (succinate dehydrogenase) or *KLLN* (killin) mutations.¹⁹ Mucocutaneous features of trichilemmomas, oral papillomatosis, and feet or hand keratosis are present in 80% of patients with PHTS, whereas colorectal polyps are found in 35 to 65% of patients.²⁰ CRC risk in PHTS is significantly increased relative to the general population, with a 4- to 10-fold increased risk²¹ or an 18% risk of CRC at the age of 60 years.²² Recommendations are for colonoscopy beginning at the age of 40 years and biannual surveillance with surgical intervention based on disease burden.¹⁸

Serrated Polyp Syndromes

Of all CRCs, 15 to 30% arise from serrated polyps rather than from adenomatous polyps. Serrated polyposis syndrome (SPS) is characterized by multiple serrated polyps throughout the colon and is accompanied by an increased CRC risk.²³ Because germline mutations for SPS are still unknown, the disease is defined clinically by (1) the presence of five or more serrated polyps proximal to the sigmoid colon, of which two are more than 10 mm in diameter; (2) the presence of one serrated polyp proximal to the sigmoid and a first-degree relative with SPS; or (3) 20 serrated polyps located throughout the colorectum, or a combination of these clinical findings. CRC risk for patients at their first SPS presentation is reported between 29 and 50%; however, this range likely overestimates the risk because of selection bias and nonstructured surveillance.²⁴ In SPS, CRC is diagnosed at a median age of 60 years.²⁴ Screening is recommended to begin at the age of 40 years and to continue at a rate of every 1 to 3 years.¹⁸

Screening and Surveillance

The American College of Gastroenterology has devised recommendations for screening and surveillance for the polyposis syndromes, outlined in ►Table 1.¹⁸

Indications and Overview of Surgical Intervention

Indications

Pediatric indications for colorectal surgery in any of the polyposis syndromes include documented or suspected

Table 1 American College of Gastroenterology Recommendations for age to begin screening and repeat screening interval for pediatric polyposis syndromes¹⁸

Polyposis syndrome	Age to begin screening, y	Repeat screening interval, y
FAP	10–15	1–2
Attenuated FAP	18–20	1–2
<i>MUTYH</i> -associated polyposis	25–30	1–2
Peutz-Jeghers syndrome	8–18	3
Juvenile polyposis syndrome	12–15	1–3
<i>PTEN</i> hamartomatous tumor syndromes	15	2
Serrated polyposis syndrome	40	2

Abbreviation: FAP, familial adenomatous polyposis.

cancer and symptoms refractory to endoscopic management. Relative indications include the presence of multiple large adenomas, a notable increase in number of adenomas, high-grade dysplasia, or the inability to complete colon surveillance because of multiple diminutive polyps. Deep family concern, particularly when there has been a death due to CRC, may prompt requests for earlier colectomy in otherwise asymptomatic children with a high volume of disease. Planning for major prophylactic surgery must consider the child's input, growth, development, and school life. Colectomy before the teenage years is reasonable in the subset of patients with thousands of polyps and a family history of early-onset cancer.²⁵

Surgical management for FAP most often includes a restorative proctocolectomy with ileal pouch anal anastomosis (IPAA).²⁶ Patients with AFAP may be successfully treated with repeated endoscopic polypectomies and may never need a colectomy, depending on ability to survey the colon and achieve polyp clearance. In addition, some patients with AFAP and *MUTYH*-associated polyposis have relative rectal sparing, allowing ileorectal anastomosis (IRA) rather than IPAA.⁵ Both groups need endoscopic monitoring for malignancy within the remaining rectum or at the anal transition zone.²⁷

For PJS, endoscopic removal of all polyps is first attempted, but colectomy is sometimes indicated when colonoscopic management is not possible or neoplasia is seen in the colonic polyps.²⁸ For JPS, colectomy may be indicated when cancer or high-grade dysplasia is detected or when the polyps cannot be adequately controlled endoscopically.²⁹ In addition, abundant polyp burden can be associated with protein-losing enteropathy, diarrhea, and electrolyte imbalances.

For SPS, development of cancer or high-grade dysplasia or the failure to endoscopically control the serrated polyp growth are indications for surgery. When indicated, prophylactic or therapeutic colectomy is with either subtotal colectomy and IRA or proctocolectomy with IPAA.³⁰

Surgical Approach

Prevention of cancer in children with polyposis syndromes is managed most effectively through total colectomy with IRA or by proctocolectomy with IPAA.³¹ Proctocolectomy with IPAA is a restorative procedure that minimizes the risk of CRC by removing almost all at-risk mucosa while maintaining reasonable bowel continence.³² This technique has been widely used in adults since the 1980s but only recently in children.³² Operative steps include proctocolectomy, endoanal mucosectomy, and IPAA with or without a diverting ileostomy. Originally an open procedure, its use progressed through laparoscopy-assisted technique, and now it is commonly performed laparoscopically.^{33,34} It has also been done successfully through a single laparoscopic incision.^{35,36}

IRA versus IPAA

A spectrum of opinion still exists among pediatric and colorectal surgeons on the optimal extent of resection for prophylactic surgery in FAP—namely, whether IRA or IPAA is the more suitable prophylactic procedure.³⁷ IRA was previously considered less invasive, involving less pelvic dissection and associated morbidity and with better functional outcomes than IPAA in selected subsets of patients.³⁸ Patients with FAP who underwent IRA rather than IPAA had reduced daily bowel frequency, less nocturnal stooling, and fewer 30-day reinterventions.³⁷ The clear disadvantages of IRA are the need for ongoing surveillance secondary to the risk of CRC in the rectum,^{39,40} leading to a secondary proctectomy (42% at 20 years).^{40,41} IPAA leaves less mucosa with the potential for dysplasia, which may be further reduced when mucosectomy is performed to undertake a hand-sewn pouch anal anastomosis.^{42,43} However, some evidence shows that mucosectomy itself can lead to a worse functional outcome and may carry the risk of leaving islands of mucosa on the muscle cuff hidden from endoscopic surveillance.^{43–45}

In some published literature, IRA has been advocated for any patients younger than 30 years, with subsequent conversion to IPAA if indicated after follow-up surveillance.⁴⁶ The purported advantage of this approach includes a less complex procedure without the need for a pouch and, importantly for children, a lower risk of infertility and impotence.^{47,48} Although IRA is feasible for patients with attenuated FAP, the increasing risk of CRC in the remaining rectum with age is considerable and has been reported at between 29 and 35% at 25 years after the original procedure.^{38,46} In that clinical setting, secondary excision of the rectum may be necessary, as reported by Booiij et al,⁴⁹ where 7 of 34 patients had completion proctectomy and 2 patients died secondary to invasive rectal cancer.⁴⁹

Functionally, there is little choice between IRA and IPAA.⁵⁰ Newer studies indicate that laparoscopic approaches may have reduced the effect of IPAA on long-term fertility.^{51,52} Furthermore, with the greater complexity of a secondary proctectomy, it may not be possible to convert an IRA into an IPAA, and if such a patient has rectal cancer, then the patient may need a completion proctectomy with end ileostomy rather than a restorative procedure.⁴¹ Thus, despite the increased complexity and potentially increased

morbidity of a proctectomy, IPAA should currently be proposed to any high-risk FAP patient as the operation of choice for optimal control of CRC risk.

Straight versus Pouch Reconstruction

Straight ileoanal anastomoses were initially championed for children because they were thought to be associated with less risk of strictures, less fecal stasis, and less inflammation of the neorectum.⁵³ They were already used commonly in pediatric surgery techniques involving pull-through for colonic aganglionosis or anorectal malformations. The straight pull-through technique retains the functional direction of peristaltic contractions and can generate physiological spike waves down to the anal anastomosis. The procedure also can be technically simpler than the pouch creation.

Although the high adaptability of children and excellent continence mechanisms were expected to mitigate the problems of high stool frequency,⁵³ creation of an ileal reservoir was seen to result in superior daily continence and less frequent daily bowel movements compared with the straight pull-through.⁵⁴ Ileal pouches are now standard of care for most pediatric reconstructive procedures.⁵⁵ Four pouch configurations have been used: lateral, S-shaped, J-shaped, and W-shaped pouches.

The J pouch configuration is favored because of its technical simplicity, quick creation, ease of the fit of the pouch into the sacrum, excellent functional outcomes, and a dearth of long-term complications.²⁶ Both the S and W pouches are hand-sutured, leading to increased complexity and a longer operating time. The S and W pouches are more vulnerable to outlet obstruction and the need for an irrigating catheter.⁵⁶ The lateral reservoir has been infrequently done in recent years, but it may be considered to reduce the tension at the anastomosis in patients with a short ileal mesentery.⁵⁷ The first IPAA described by Park and Nicholls³² involved an S-shaped ileal pouch mobilized through a mucosectomized rectum and hand-sewn to the dentate line. The J pouch, initially described by Utsunomiya and colleagues,⁵⁸ has now become the most widely used, being simpler than other techniques, highly functional, and less prone to long-term complications.⁵⁵

To construct the J pouch, the ileal mesentery should be mobilized adjacent to the superior mesenteric vessels in a length sufficient to allow the ileum to be anastomosed to the anus without tension. To create the pouch, the ileum is folded back on itself for a pouch length of 10 to 12 cm, and a stapling device is used to form the anastomosis. Sutures also are placed in a seromuscular manner at the top of the suture line, to divert tension off the area. The length of the J pouch was shortened after the realization that longer pouches led to increased stool stasis and pouchitis.⁵⁸

Straight versus J-Shaped Pouch IPAA

A pediatric meta-analysis comparing the straight IAA versus the J pouch IAA indicated a trend for lower daily stool counts in children who had a pouch rather than a straight pull-through.⁵⁹ However, a further multicenter study indicated that although straight anastomosis had higher stool frequency and the J pouch had increased pouchitis at 12 months from surgery,

these problems regressed over the next 12 months from surgery and functional stooling scores became similar between the two groups.⁵⁵ The patients with J-style pouch anastomoses maintained lower stool frequency and marginally better continence; yet, these differences became small later and may have had minimal clinical significance. Continence stayed excellent regardless of which technique was used (92 vs. 97% at 1 year).⁵⁵ Straight anastomosis reduced the risk of pouchitis associated with an ileal reservoir. However, rates of pouchitis were lower in anastomosis undertaken for FAP than for ulcerative colitis (UC), and that incidence itself decreased after 24 months.⁵⁵ Pull-through failures requiring reversion to a permanent ileostomy occurred in 6% of patients, and that percentage did not vary with any statistical significance between groups (8% S-shaped IAA [most often secondary to stooling frequency] and 5% J-shaped PAA [most often secondary to pouchitis]).⁵⁵

S-Shaped, J-Shaped, or W-Shaped Pouch

Evaluation has been extensive on the different types of ileal pouch reservoirs.^{60–63} Anatomically, a J pouch fits optimally into the sacrum, with the larger end of the J resting on the levators. An S pouch creates a longer spout, which carries the potential to lead to obstruction or kinking. A meta-analysis of 18 studies that contained 1,519 patients—689 with a J pouch, 306 with a W pouch, and 524 S pouch—was not able to detect any statistically significant difference in early postoperative complications.⁶⁰ In the short term, daily stool frequency was greater for patients with a J pouch than an S or W pouch.⁶⁰ Obstructed evacuation was more common in patients with an S- or W-shaped pouch. No long-term significant difference was detected between patients with an S-shaped pouch and those with a J-shaped one for most functional outcomes.⁶⁰ Variation within the patient subset techniques (e.g., hand-sewn vs. stapled and laparoscopic vs. open) and the patient indications (FAP vs. UC) may make it difficult to detect any true difference in outcomes for individual patients. At this time, the risk of obstruction in the S-shaped pouch is further decreased with suggestions that the exit conduit of an S-shaped pouch should be no greater than 2 to 2.5 cm in length, as opposed to 5 cm in its initial designs.⁵⁶

Hand-Sewn versus Stapled Anastomosis

Currently, no consensus exists on whether to undertake hand-sewn anastomosis or stapled anastomosis in children. The IPAA procedure was originally designed to include a rectal mucosectomy and hand-sewn ileal pouch to anal canal anastomosis. By preserving the anal canal transition zone and a cuff of distal rectum, a double-stapled technique was thought to potentially result in improved postoperative function.⁶⁴ To this day, though, controversy continues regarding which of these techniques is superior.⁶⁵

The stapled technique (as illustrated in ► Fig. 1) is thought to be technically easier and thus has been proposed to be less prone to leak or stenosis complications and to have fewer sepsis-related pouch excisions.⁶⁶ However, in patients with FAP, a matched-pair analysis indicated a nonstatistically significant tendency toward a higher rate of early postoperative complications—in particular, stenosis—after stapled IPAA.⁶⁷




Fig. 1 Stapled ileoanal anastomosis. (A) The anastomosis is undertaken with the circular stapler as indicated. (B) Shorter cuff lengths are possible with this technique (©Mayo Clinic).

Pouchitis outcomes are seen as comparable between hand-sewn and stapled anastomosis.⁶⁸ The stapled technique tends to be undertaken in larger children, in keeping with the adult colorectal experience and with the availability of appropriate sizes of staplers. Stapled anastomosis leads to a longer cuff length, which was hypothesized to improve functional outcomes. However, a meta-analysis of prospective studies did not detect a statistically significant difference in long-term functional outcomes between hand-sewn or stapled anastomosis.⁶⁹

For the UC patient population, recent literature supports a stapled IPAA, with improved functional results and fewer postoperative complications than the hand-sewn anastomosis.^{66,68,70} A mucosectomy and hand-sewn IPAA are entirely necessary in certain situations, though: neoplasia in the anal transition zone, large polyp burden in the low rectum, a revision IPAA, and small patient size in a younger patient for whom stapling devices do not fit.⁴³

With or without a Diverting Stoma

The restorative proctocolectomy can be performed with or without a diverting loop ileostomy.^{25,71} It was initially a standard practice to perform IPAA as two- or three-stage procedures, in particular using a diverting loop ileostomy to reduce the risk or impact of pelvic sepsis,⁷² which is known to be a critical determinant of pouch failure.⁷³ A North American survey indicated that most colorectal surgeons believed that proximal diversion at the time of IPAA was still indicated.⁷⁴ In particular, there is concern that a pelvic complication could lead to pouch failure and create the need for a permanent ileostomy.^{75–77} Several studies have shown a low complication rate even with a single-stage approach.^{71,78} As such, the incidence of one- and two-stage IPAA has increased over time and are considered in younger patients particularly.²⁵ Evaluation of the Mayo Clinic pediatric experience indicated that patients with one-stage procedures had higher rates of leak and higher need for reoperation than patients with two-stage IPAA.²⁵ The

patients had better long-term bowel function with a one-stage IPAA.²⁵ Hence, current recommendations cite a diverting ileostomy as the standard of care, with one-stage procedures only in highly selected patients.

Anatomical Reach

Regardless of which pouch technique is chosen, the anastomosis must reach the anus without undue tension. In univariate and multivariate analyses, Heuschen et al⁷⁹ showed that anastomotic tension was a significant risk factor for early complications in patients with FAP, in particular for pouch-related septic complications. For a pouch, the site of ileal reach needs to be proximal to the terminal staple line. For instance, for the J-shaped pouch, a site 16 to 20 cm proximal to the terminal ileal staple line is the actual anastomotic ileoanal apex. Therefore, the reach needs to be longer than for a straight pull-through procedure.

The pediatric mesentery is not necessarily more flexible than the adult mesentery; so, mesenteric windows and lengthening maneuvers are necessary to minimize tension on the anastomosis (→ **Fig. 2C–E**). Relaxing incisions in the peritoneum can be made both front and back. Typically, approximately five relaxing incisions are made, and these will likely gain up to 2 cm in length.⁸⁰

The main feature limiting reach is the mesenteric vessels, some of which need to be sacrificed to optimize reach and minimize tension (→ **Fig. 2A**). Initial studies by Smith et al⁸¹ identified key principles that should be observed during the management of the mesenteric circulation. By dividing the distal superior mesenteric artery but preserving the ileocolic artery, an extra 2.5 cm in length on average could be generated. Martel et al^{82,83} analyzed whether a high division of the superior mesenteric artery with preservation of the ileocolic artery was safe. They showed that this approach led to an increased reach of 3.5 cm on average and often necessitated a longer ileal resection. Burnstein et al⁸⁰ stated that division of




Fig. 2 Maneuvers to increase mesenteric reach in ileal pouch anal anastomosis. (A) Initial mobilization of the ileal mesentery away from the retroperitoneum by optimal dissection to the mesenteric root of the SMA (IC artery, ileocolic artery; RC, right colic artery; -MC, middle colic artery). (B) This exposes the aorta (ao), duodenum (D), right ureter (Ur), and gonadal vessels (GV). (C) The mesenteric reach can be further extended by scoring the peritoneum stepwise. (D and E) Ligation of the ileocolic with preservation of the distal SMA can provide further reach for the ileal J pouch (©Mayo Clinic).

either of the major mesenteric vessels was usually unnecessary, particularly in the pediatric population. Their technique, instead, involved division of two or three smaller ileal mesenteric arteries between the primary and secondary arcades, resulting in an average of 2 to 5 cm of extra length.⁸⁰ An alternative technique if reach is a more challenging involved preservation of the right branch of the middle colic and the marginal artery of the right colon with division of the right colic and ileocolic arteries at their origin. An average extra 11 cm was gained through this technique.⁸⁴ Cherqui et al⁸⁵ showed that mobilization to the mesentery root added 1 cm in some patients⁸⁵ (►Fig. 2B). These techniques can be used in combination with optimized reach, depending on the patient's anatomy.⁸⁶

Laparoscopic versus Open Approach

Minimally invasive laparoscopic approaches for these young and asymptomatic patients are appealing. All currently published reports indicate that laparoscopic restorative proctoco-

lectomy is safe and may result in a shorter length of hospital stay and a better cosmetic outcome.³³ The drawbacks that investigators observed include longer operating times initially and a steep learning curve. With technique improvement, better outcomes, including reduced adhesions, less postoperative pain, and ileus, and a quicker return to normal activities may emerge for laparoscopic IPAA.⁸⁷⁻⁸⁹ In addition, single-port laparoscopic approaches are being innovated to further advance minimally invasive management for this young patient population.^{35,36}

Contraindications to IPAA

Absolute contraindications to IPAA are few. Occasionally, when the mesentery is foreshortened, a restorative surgery is unable to proceed despite mesenteric lengthening procedures. This result is particularly the case for patients with a thick mesentery and a high body mass index. Malrotation, recurrent desmoid tumors, and prior hepatoblastoma surgery are all relative

contraindications to proceeding to IPAA but not to proceeding to colectomy and end ileostomy. Similarly, rectal dysplasia mandates a full proctectomy with mesorectal excision rather than an IPAA.⁹⁰ Children or young adults with rectal dysplasia or malignancy will reasonably resist permanent ileostomy as the only option, but the procedure should not be delayed when indications to operate are strong. Judgment is of course paramount. A patient at high risk for desmoid tumors should not undergo early prophylactic IPAA. However, with the nearly 100% incidence of invasive adenocarcinoma by the age of 50 years, concerns for desmoid complications, while important, are outweighed by the cancer risk in otherwise healthy patients.

The decision to operate continues to be based on cancer prophylaxis. When familial patterns and genotype identify high-risk patients and surveillance confirms imminent malignancy, colectomy should proceed despite the relative contraindications of a permanent ileostomy.

IPAA Morbidity and Long-Term Outcomes

Overall, morbidity from prophylactic IPAA is substantial and ranges from 10 to 25% for patients with FAP—notably better than for patients undergoing surgery for UC.^{91,92} Anastomotic leak and sepsis range from 0 to 9%; the main risk factor is anastomotic tension.⁷⁹ Nonfibrotic strictures at the anastomosis are common, but they respond well with anal dilatation. Fibrotic strictures may occur secondary to intraoperative complications and may more often need surgical intervention.⁹³ Pouch failure requiring excision (risk, 4–8%) is associated with chronic pouchitis and with pelvic leak.⁵⁵ Revision IPAA is usually possible with an acceptable outcome.^{57,94} Small-bowel obstruction rate is high after IPAA; MacLean et al⁹⁵ reported a cumulative risk of 31% and need for surgical intervention in 7.5%. Pouchitis after IPAA is seen less often in patients with FAP than in patients with UC (0–11%).^{79,96}

Stool frequency has been reported as four to six bowel movements daily, and average episodes of nighttime stooling were zero to one bowel movement.^{92,97,98} Rates of normal daytime continence were impressively high at 80 to 95% of patients. Nighttime fecal spotting rates were moderate overall at 32 to 42%, and approximately 1% of patients reported some soiling at night.^{92,98–101} Long-term follow-up in large patient populations has indicated that continence does not deteriorate significantly over time.^{99,101} Pouchitis prevalence following IPAA for FAP was low (<10%).⁹⁶

An important concern for functional outcomes is postoperative fertility, which decreases to 54% after IPAA in FAP patients.¹⁰² Pediatric patients with FAP are often childless at the time of surgical intervention, and the desire for future fertility may be substantial.¹⁰³ IPAA does not impair childbirth or even pregnancy,¹⁰⁴ but it does have a considerable effect on female fertility,^{102,105} as well as a small risk of impairment in male fertility.¹⁰⁶ A close rectal wall proctectomy technique may reduce the effect on future fertility.¹⁰⁴ Experts hypothesize that pelvic adhesions from the pelvic dissection may be responsible.^{107,108} A subset of patients who underwent colectomy with either IPAA or terminal ileostomy underwent hysterosalpingography analysis. The imaging showed unilateral or bilateral fallopian tube

obstruction in 52% of patients.^{109,110} Newer studies indicate that laparoscopic approaches may significantly reduce the effect of the operation on a patient's long-term fertility.^{51,52}

Pregnancies are most often uneventful in patients who underwent IPAA; most pregnancies lead to term labor and delivery without complications. Neither the pregnancy nor the labor adversely affects pouch function, and there is no mandate for a Cesarean section.^{70,104,111}

Urinary and sexual dysfunction is possible after IPAA, and the incidence varies depending on the rectal dissection technique.¹⁰⁴ With a close rectal wall dissection, minimal impact on urinary and sexual dysfunction was seen.^{26,109} Colwell and Gray¹⁰⁶ reported rates of 0.5 to 1.5% for erectile dysfunction and 3 to 4% for ejaculatory dysfunction. Incidence of female sexual dysfunction after IPAA has been less thoroughly analyzed, but the same rates of 3 to 22% dyspareunia are reported.¹¹² In addition, concerns for stool leakage have led to inhibited sexual interactions in 3% of female patients after IPAA.¹⁰⁶

Almost all quality-of-life outcomes after IPAA were comparable to those of healthy children.^{97,99,100} In a Cleveland Clinic series, overall long-term quality of life after IPAA was rated as good to excellent in 99% of FAP patients.¹⁰¹ Patient satisfaction was also markedly high after IPAA.^{97,99,101} Daily activities—from social, home, travel, sports, and sexual—are affected minimally in the long term, with only 2% of patients reporting adverse outcomes in these activity areas.¹¹³ In the pediatric setting, quality of life, physical functions, mental health, and self-esteem were all seen to be equivalent to healthy children.⁹⁷ Parental concerns about the health of the child continued to persist, however, even in the absence of physical or functional abnormalities.⁹⁷

Conclusion

Inherited polyposis syndromes present a distinctive subset of patients whose risk of CRC may be obviated by prophylactic resection of all colorectal mucosa and IPAA reconstruction.¹¹⁴ Diagnosis is becoming earlier with increasing awareness and understanding, and hence these patients are increasingly offered surgical interventions during their younger years. IPAA should be offered to the majority of patients with polyposis syndromes when their risks of colorectal malignancy have been established after serial endoscopic surveillance. The procedure is safe, results in removal of all colorectal mucosa and provides CRC risk reduction, offers good functional outcomes, and has limited complications. Continued technical advances and improved surgical techniques will likely further improve the functional outcome and patient satisfaction after IPAA.

References

- 1 Burn J, Chapman P, Delhanty J, et al. The UK Northern region genetic register for familial adenomatous polyposis coli: use of age of onset, congenital hypertrophy of the retinal pigment epithelium, and DNA markers in risk calculations. *J Med Genet* 1991;28(05):289–296

- 2 Kerr SE, Thomas CB, Thibodeau SN, Ferber MJ, Halling KC. APC germline mutations in individuals being evaluated for familial adenomatous polyposis: a review of the Mayo Clinic experience with 1591 consecutive tests. *J Mol Diagn* 2013;15(01):31–43
- 3 Bussey H. *Familial Polyposis Coli: Family Studies, Histopathology, Differential Diagnosis, and Results of Treatment*. Baltimore, MD: Johns Hopkins University Press; 1975
- 4 Vasen HF, Möslin G, Alonso A, et al. Guidelines for the clinical management of familial adenomatous polyposis (FAP). *Gut* 2008;57(05):704–713
- 5 Burt RW, Leppert MF, Slattery ML, et al. Genetic testing and phenotype in a large kindred with attenuated familial adenomatous polyposis. *Gastroenterology* 2004;127(02):444–451
- 6 Buecher B. Colorectal adenomatous polyposis syndromes: genetic determinism, clinical presentation and recommendations for care. *Bull Cancer* 2016;103(02):199–209
- 7 Palles C, Cazier JB, Howarth KM, et al; CORGI Consortium; WGS500 Consortium. Germline mutations affecting the proof-reading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet* 2013;45(02):136–144
- 8 Bellido F, Pineda M, Aiza G, et al. POLE and POLD1 mutations in 529 kindred with familial colorectal cancer and/or polyposis: review of reported cases and recommendations for genetic testing and surveillance. *Genet Med* 2016;18(04):325–332
- 9 Spier I, Drichel D, Kerick M, et al. Low-level APC mutational mosaicism is the underlying cause in a substantial fraction of unexplained colorectal adenomatous polyposis cases. *J Med Genet* 2016;53(03):172–179
- 10 Chubb D, Broderick P, Frampton M, et al. Genetic diagnosis of high-penetrance susceptibility for colorectal cancer (CRC) is achievable for a high proportion of familial CRC by exome sequencing. *J Clin Oncol* 2015;33(05):426–432
- 11 Tomlinson IP, Houlston RS. Peutz-Jeghers syndrome. *J Med Genet* 1997;34(12):1007–1011
- 12 McGrath DR, Spigelman AD. Preventive measures in Peutz-Jeghers syndrome. *Fam Cancer* 2001;1(02):121–125
- 13 Hearle N, Schumacher V, Menko FH, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. *Clin Cancer Res* 2006;12(10):3209–3215
- 14 Wirtzfeld DA, Petrelli NJ, Rodriguez-Bigas MA. Hamartomatous polyposis syndromes: molecular genetics, neoplastic risk, and surveillance recommendations. *Ann Surg Oncol* 2001;8(04):319–327
- 15 Chow E, Macrae F. A review of juvenile polyposis syndrome. *J Gastroenterol Hepatol* 2005;20(11):1634–1640
- 16 Schreiber IR, Baker M, Amos C, McGarrity TJ. The hamartomatous polyposis syndromes: a clinical and molecular review. *Am J Gastroenterol* 2005;100(02):476–490
- 17 Cichy W, Klineciewicz B, Plawski A. Juvenile polyposis syndrome. *Arch Med Sci* 2014;10(03):570–577
- 18 Syngal S, Brand RE, Church JM, Giardiello FM, Hampel HL, Burt RW; American College of Gastroenterology. ACG clinical guideline: genetic testing and management of hereditary gastrointestinal cancer syndromes. *Am J Gastroenterol* 2015;110(02):223–262, quiz 263
- 19 Ngeow J, Mester J, Rybicki LA, Ni Y, Milas M, Eng C. Incidence and clinical characteristics of thyroid cancer in prospective series of individuals with Cowden and Cowden-like syndrome characterized by germline PTEN, SDH, or KLLN alterations. *J Clin Endocrinol Metab* 2011;96(12):E2063–E2071
- 20 Omundsen M, Lam FF. The other colonic polyposis syndromes. *ANZ J Surg* 2012;82(10):675–681
- 21 Tan MH, Mester JL, Ngeow J, Rybicki LA, Orloff MS, Eng C. Lifetime cancer risks in individuals with germline PTEN mutations. *Clin Cancer Res* 2012;18(02):400–407
- 22 Nieuwenhuis MH, Kets CM, Murphy-Ryan M, et al. Is colorectal surveillance indicated in patients with PTEN mutations? *Colorectal Dis* 2012;14(09):e562–e566
- 23 Snover DC, Ahnen DJ, Burt RW, et al. Serrated polyps of the colon and rectum and serrated polyposis. In: Bosman T, Carneiro F, Hruban R, et al. eds. *WHO Classification of Tumours of the Digestive System*. Lyon2010:160–165
- 24 Ijspeert JE, Rana SA, Atkinson NS, et al; Dutch Workgroup Serrated Polyps & Polyposis (WASP). Clinical risk factors of colorectal cancer in patients with serrated polyposis syndrome: a multicentre cohort analysis. *Gut* 2017;66(2):278–284
- 25 Kennedy RD, Potter DD, Moir CR, El-Youssef M. The natural history of familial adenomatous polyposis syndrome: a 24 year review of a single center experience in screening, diagnosis, and outcomes. *J Pediatr Surg* 2014;49(01):82–86
- 26 Kartheuser A, Stangherlin P, Brandt D, Remue C, Sempoux C. Restorative proctocolectomy and ileal pouch-anal anastomosis for familial adenomatous polyposis revisited. *Fam Cancer* 2006;5(03):241–260, discussion 261–262
- 27 Ooi BS, Remzi FH, Gramlich T, Church JM, Preen M, Fazio VW. Anal transitional zone cancer after restorative proctocolectomy and ileoanal anastomosis in familial adenomatous polyposis: report of two cases. *Dis Colon Rectum* 2003;46(10):1418–1423, discussion 1422–1423
- 28 Giardiello FM, Trimath JD. Peutz-Jeghers syndrome and management recommendations. *Clin Gastroenterol Hepatol* 2006;4(04):408–415
- 29 Oncel M, Church JM, Remzi FH, Fazio VW. Colonic surgery in patients with juvenile polyposis syndrome: a case series. *Dis Colon Rectum* 2005;48(01):49–55, discussion 55–56
- 30 Young JP, Parry S. Risk factors: hyperplastic polyposis syndrome and risk of colorectal cancer. *Nat Rev Gastroenterol Hepatol* 2010;7(11):594–595
- 31 Church J, Burke C, McGannon E, Pastean O, Clark B. Risk of rectal cancer in patients after colectomy and ileorectal anastomosis for familial adenomatous polyposis: a function of available surgical options. *Dis Colon Rectum* 2003;46(09):1175–1181
- 32 Parks AG, Nicholls RJ. Proctocolectomy without ileostomy for ulcerative colitis. *BMJ* 1978;2(6130):85–88
- 33 Larson DW, Cima RR, Dozois EJ, et al. Safety, feasibility, and short-term outcomes of laparoscopic ileal-pouch-anal anastomosis: a single institutional case-matched experience. *Ann Surg* 2006;243(05):667–670, discussion 670–672
- 34 Fraser JD, Garey CL, Laituri CA, Sharp RJ, Ostlie DJ, St Peter SD. Outcomes of laparoscopic and open total colectomy in the pediatric population. *J Laparoendosc Adv Surg Tech A* 2010;20(07):659–660
- 35 Potter DD, Tung J, Faubion WA Jr, Moir C. Single-incision laparoscopic colon and rectal surgery for pediatric inflammatory bowel disease and polyposis syndromes. *J Laparoendosc Adv Surg Tech A* 2012;22(02):203–207
- 36 Schlager A, Siddharthan RV, Santore MT, Balci O, Clifton MS, Heiss KF. Single-incision total proctocolectomy and ileal pouch anal anastomosis in pediatric patients: lessons learned. *J Laparoendosc Adv Surg Tech A* 2015;25(09):782–785
- 37 Aziz O, Athanasiou T, Fazio VW, et al. Meta-analysis of observational studies of ileorectal versus ileal pouch-anal anastomosis for familial adenomatous polyposis. *Br J Surg* 2006;93(04):407–417
- 38 Bülow C, Vasen H, Järvinen H, Björk J, Bisgaard ML, Bülow S. Ileorectal anastomosis is appropriate for a subset of patients with familial adenomatous polyposis. *Gastroenterology* 2000;119(06):1454–1460
- 39 DeCosse JJ. Surgical prophylaxis of familial colon cancer: prevention of death from familial colorectal cancer. *J Natl Cancer Inst Monogr* 1995;(17):31–32
- 40 Smith JC, Schäffer MW, Ballard BR, et al. Adenocarcinomas after prophylactic surgery for familial adenomatous polyposis. *J Cancer Ther* 2013;4(01):260–270
- 41 von Roon AC, Tekkis PP, Lovegrove RE, Neale KF, Phillips RK, Clark SK. Comparison of outcomes of ileal pouch-anal anastomosis for

- familial adenomatous polyposis with and without previous ileorectal anastomosis. *Br J Surg* 2008;95(04):494–498
- 42 von Roon AC, Will OC, Man RF, et al. Mucosectomy with hand-sewn anastomosis reduces the risk of adenoma formation in the anorectal segment after restorative proctocolectomy for familial adenomatous polyposis. *Ann Surg* 2011;253(02):314–317
- 43 Remzi FH, Church JM, Bast J, et al. Mucosectomy vs. stapled ileal pouch-anal anastomosis in patients with familial adenomatous polyposis: functional outcome and neoplasia control. *Dis Colon Rectum* 2001;44(11):1590–1596
- 44 Chambers WM, McC Mortensen NJ. Should ileal pouch-anal anastomosis include mucosectomy? *Colorectal Dis* 2007;9(05):384–392
- 45 M'Koma AE, Moses HL, Adunyah SE. Inflammatory bowel disease-associated colorectal cancer: proctocolectomy and mucosectomy do not necessarily eliminate pouch-related cancer incidences. *Int J Colorectal Dis* 2011;26(05):533–552
- 46 Nugent KP, Phillips RK. Rectal cancer risk in older patients with familial adenomatous polyposis and an ileorectal anastomosis: a cause for concern. *Br J Surg* 1992;79(11):1204–1206
- 47 Harnoy Y, Desfourneaux V, Bouguen G, et al. Sexuality and fertility outcomes after hand sewn versus stapled ileal pouch anal anastomosis for ulcerative colitis. *J Surg Res* 2016;200(01):66–72
- 48 van Balkom KA, Beld MP, Visschers RG, van Gemert WG, Breukink SO. Long-term results after restorative proctocolectomy with ileal pouch-anal anastomosis at a young age. *Dis Colon Rectum* 2012;55(09):939–947
- 49 Booi KA, Mathus-Vliegen EM, Taminiau JA, et al. Evaluation of 28 years of surgical treatment of children and young adults with familial adenomatous polyposis. *J Pediatr Surg* 2010;45(03):525–532
- 50 van Duijvendijk P, Slors JF, Taat CW, Oosterveld P, Vasen HF. Functional outcome after colectomy and ileorectal anastomosis compared with proctocolectomy and ileal pouch-anal anastomosis in familial adenomatous polyposis. *Ann Surg* 1999;230(05):648–654
- 51 Bartels SA, D'Hoore A, Cuesta MA, Bendsdorp AJ, Lucas C, Bemelman WA. Significantly increased pregnancy rates after laparoscopic restorative proctocolectomy: a cross-sectional study. *Ann Surg* 2012;256(06):1045–1048
- 52 Beyer-Berjot L, Maggiori L, Birnbaum D, Lefevre JH, Berdah S, Panis Y. A total laparoscopic approach reduces the infertility rate after ileal pouch-anal anastomosis: a 2-center study. *Ann Surg* 2013;258(02):275–282
- 53 Coran AG. A personal experience with 100 consecutive total colectomies and straight ileoanal endorectal pull-throughs for benign disease of the colon and rectum in children and adults. *Ann Surg* 1990;212(03):242–247, discussion 247–248
- 54 Rintala RJ, Lindahl H. Restorative proctocolectomy for ulcerative colitis in children—is the J-pouch better than straight pull-through? *J Pediatr Surg* 1996;31(04):530–533
- 55 Seetharamaiah R, West BT, Ignash SJ, et al. Outcomes in pediatric patients undergoing straight vs J pouch ileoanal anastomosis: a multicenter analysis. *J Pediatr Surg* 2009;44(07):1410–1417
- 56 Wu XR, Kirat HT, Kalady MF, Church JM; Wu XR1. Restorative proctocolectomy with a handsewn IPAA: S-pouch or J-pouch? *Dis Colon Rectum* 2015;58(02):205–213
- 57 Fonkalsrud EW, Bustorff-Silva J. Reconstruction for chronic dysfunction of ileoanal pouches. *Ann Surg* 1999;229(02):197–204
- 58 Fujita S, Kusunoki M, Shoji Y, Owada T, Utsunomiya J. Quality of life after total proctocolectomy and ileal J-pouch-anal anastomosis. *Dis Colon Rectum* 1992;35(11):1030–1039
- 59 Tilney HS, Constantinides V, Ioannides AS, Tekkis PP, Darzi AW, Haddad MJ. Pouch-anal anastomosis vs straight ileoanal anastomosis in pediatric patients: a meta-analysis. *J Pediatr Surg* 2006;41(11):1799–1808
- 60 Lovegrove RE, Heriot AG, Constantinides V, et al. Meta-analysis of short-term and long-term outcomes of J, W and S ileal reservoirs for restorative proctocolectomy. *Colorectal Dis* 2007;9(04):310–320
- 61 McHugh SM, Diamant NE, McLeod R, Cohen Z. S-pouches vs. J-pouches. A comparison of functional outcomes. *Dis Colon Rectum* 1987;30(09):671–677
- 62 Tuckson WB, Fazio VW. Functional comparison between double and triple ileal loop pouches. *Dis Colon Rectum* 1991;34(01):17–21
- 63 Gemlo BT, Belmonte C, Wiltz O, Madoff RD. Functional assessment of ileal pouch-anal anastomotic techniques. *Am J Surg* 1995;169(01):137–141, discussion 141–142
- 64 Geiger JD, Teitelbaum DH, Hirschl RB, Coran AG. A new operative technique for restorative proctocolectomy: the endorectal pull-through combined with a double-stapled ileo-anal anastomosis. *Surgery* 2003;134(03):492–495
- 65 Kirat HT, Remzi FH, Kiran RP, Fazio VW. Comparison of outcomes after hand-sewn versus stapled ileal pouch-anal anastomosis in 3,109 patients. *Surgery* 2009;146(04):723–729, discussion 729–730
- 66 Ziv Y, Fazio VW, Church JM, Lavery IC, King TM, Ambrosetti P. Stapled ileal pouch anal anastomoses are safer than handsewn anastomoses in patients with ulcerative colitis. *Am J Surg* 1996;171(03):320–323
- 67 Ganschow P, Warth R, Hinz U, Büchler MW, Kadmon M. Early postoperative complications after stapled vs handsewn restorative proctocolectomy with ileal pouch-anal anastomosis in 148 patients with familial adenomatous polyposis coli: a matched-pair analysis. *Colorectal Dis* 2014;16(02):116–122
- 68 McIntyre PB, Pemberton JH, Beart RW Jr, Devine RM, Nivatvongs S. Double-stapled vs. handsewn ileal pouch-anal anastomosis in patients with chronic ulcerative colitis. *Dis Colon Rectum* 1994;37(05):430–433
- 69 Schluender SJ, Mei L, Yang H, Fleshner PR. Can a meta-analysis answer the question: is mucosectomy and handsewn or double-stapled anastomosis better in ileal pouch-anal anastomosis? *Am Surg* 2006;72(10):912–916
- 70 Farouk R, Pemberton JH, Wolff BG, Dozois RR, Browning S, Larson D. Functional outcomes after ileal pouch-anal anastomosis for chronic ulcerative colitis. *Ann Surg* 2000;231(06):919–926
- 71 Ryan DP, Doody DP. Restorative proctocolectomy with and without protective ileostomy in a pediatric population. *J Pediatr Surg* 2011;46(01):200–203
- 72 Martin LW, LeCoultré C, Schubert WK. Total colectomy and mucosal proctectomy with preservation of continence in ulcerative colitis. *Ann Surg* 1977;186(04):477–480
- 73 Hahnloser D, Pemberton JH, Wolff BG, Larson DR, Crownhart BS, Dozois RR. Results at up to 20 years after ileal pouch-anal anastomosis for chronic ulcerative colitis. *Br J Surg* 2007;94(03):333–340
- 74 de Montbrun SL, Johnson PM. Proximal diversion at the time of ileal pouch-anal anastomosis for ulcerative colitis: current practices of North American colorectal surgeons. *Dis Colon Rectum* 2009;52(06):1178–1183
- 75 Galandiuk S, Scott NA, Dozois RR, et al. Ileal pouch-anal anastomosis. Reoperation for pouch-related complications. *Ann Surg* 1990;212(04):446–452, discussion 452–454
- 76 Baixauli J, Delaney CP, Wu JS, Remzi FH, Lavery IC, Fazio VW. Functional outcome and quality of life after repeat ileal pouch-anal anastomosis for complications of ileoanal surgery. *Dis Colon Rectum* 2004;47(01):2–11
- 77 Lillehei CW, Leichtner A, Bousvaros A, Shamberger RC. Restorative proctocolectomy and ileal pouch-anal anastomosis in children. *Dis Colon Rectum* 2009;52(09):1645–1649
- 78 Sugerman HJ, Sugerman EL, Meador JG, Newsome HH Jr, Kellum JM Jr, DeMaria EJ. Ileal pouch anal anastomosis without ileal diversion. *Ann Surg* 2000;232(04):530–541

- 79 Heuschen UA, Hinz U, Allemeyer EH, et al. Risk factors for ileoanal J pouch-related septic complications in ulcerative colitis and familial adenomatous polyposis. *Ann Surg* 2002;235(02):207–216
- 80 Burnstein MJ, Schoetz DJ Jr, Collier JA, Veidenheimer MC. Technique of mesenteric lengthening in ileal reservoir-anal anastomosis. *Dis Colon Rectum* 1987;30(11):863–866
- 81 Smith L, Friend WG, Medwell SJ. The superior mesenteric artery. The critical factor in the pouch pull-through procedure. *Dis Colon Rectum* 1984;27(11):741–744
- 82 Martel P, Blanc P, Bothereau H, Malafosse M, Gallot D. Comparative anatomical study of division of the ileocolic pedicle or the superior mesenteric pedicle for mesenteric lengthening. *Br J Surg* 2002;89(06):775–778
- 83 Martel P, Majery N, Savigny B, Sezeur A, Gallot D, Malafosse M. Mesenteric lengthening in ileoanal pouch anastomosis for ulcerative colitis: Is high division of the superior mesenteric pedicle a safe procedure? *Dis Colon Rectum* 1998;41(07):862–866, discussion 866–867
- 84 Goes RN, Nguyen P, Huang D, Beart RWJ Jr. Lengthening of the mesentery using the marginal vascular arcade of the right colon as the blood supply to the ileal pouch. *Dis Colon Rectum* 1995;38(08):893–895
- 85 Cherqui D, Valleur P, Perniceni T, Hautefeuille P. Inferior reach of ileal reservoir in ileoanal anastomosis. Experimental anatomic and angiographic study. *Dis Colon Rectum* 1987;30(05):365–371
- 86 Chu DI, Tognelli J, Kartheuser AH, Dozois EJ. Strategy for the difficult-to-reach ileal pouch-anal anastomosis: technical steps of an in vivo application of a mesenteric-lengthening technique. *Tech Coloproctol* 2015;19(11):705–709
- 87 Garrard CL, Clements RH, Nanney L, Davidson JM, Richards WO. Adhesion formation is reduced after laparoscopic surgery. *Surg Endosc* 1999;13(01):10–13
- 88 Linden BC, Bairdain S, Zurakowski D, Shamberger RC, Lillehei CW. Comparison of laparoscopic-assisted and open total proctocolectomy and ileal pouch anal anastomosis in children and adolescents. *J Pediatr Surg* 2013;48(07):1546–1550
- 89 Causey MW, Stoddard D, Johnson EK, et al. Laparoscopy impacts outcomes favorably following colectomy for ulcerative colitis: a critical analysis of the ACS-NSQIP database. *Surg Endosc* 2013;27(02):603–609
- 90 Penna C, Turet E, Daude F, Parc R. Results of ileal J-pouch-anal anastomosis in familial adenomatous polyposis complicated by rectal carcinoma. *Dis Colon Rectum* 1994;37(02):157–160
- 91 Barton JG, Paden MA, Lane M, Postier RG. Comparison of post-operative outcomes in ulcerative colitis and familial polyposis patients after ileoanal pouch operations. *Am J Surg* 2001;182(06):616–620
- 92 Salemans JM, Nagengast FM, Lubbers EJ, Kuipers JH. Postoperative and long-term results of ileal pouch-anal anastomosis for ulcerative colitis and familial polyposis coli. *Dig Dis Sci* 1992;37(12):1882–1889
- 93 Prudhomme M, Dozois RR, Godlewski G, Mathison S, Fabbro-Peray P. Anal canal strictures after ileal pouch-anal anastomosis. *Dis Colon Rectum* 2003;46(01):20–23
- 94 Dayton MT. Redo ileal pouch-anal anastomosis for malfunctioning pouches—acceptable alternative to permanent ileostomy? *Am J Surg* 2000;180(06):561–564, discussion 565
- 95 MacLean AR, Cohen Z, MacRae HM, et al. Risk of small bowel obstruction after the ileal pouch-anal anastomosis. *Ann Surg* 2002;235(02):200–206
- 96 Kniot WA, Williams MR, Keighley MR. Pouchitis following colectomy and ileal reservoir construction for familial adenomatous polyposis. *Br J Surg* 1990;77(11):1283
- 97 Stavlo PL, Libsch KD, Rodeberg DA, Moir CR. Pediatric ileal pouch-anal anastomosis: functional outcomes and quality of life. *J Pediatr Surg* 2003;38(06):935–939
- 98 Penna C, Turet E, Kartheuser A, Hannoun L, Nordlinger B, Parc R. Function of ileal J pouch-anal anastomosis in patients with familial adenomatous polyposis. *Br J Surg* 1993;80(06):765–767
- 99 Coffey JC, Winter DC, Neary P, Murphy A, Redmond HP, Kirwan WO. Quality of life after ileal pouch-anal anastomosis: an evaluation of diet and other factors using the Cleveland Global Quality of Life instrument. *Dis Colon Rectum* 2002;45(01):30–38
- 100 Robb B, Pritts T, Gang G, et al. Quality of life in patients undergoing ileal pouch-anal anastomosis at the University of Cincinnati. *Am J Surg* 2002;183(04):353–360
- 101 Fazio VW, O'Riordain MG, Lavery IC, et al. Long-term functional outcome and quality of life after stapled restorative proctocolectomy. *Ann Surg* 1999;230(04):575–584, discussion 584–586
- 102 Olsen KØ, Joelsson M, Laurberg S, Oresland T. Fertility after ileal pouch-anal anastomosis in women with ulcerative colitis. *Br J Surg* 1999;86(04):493–495
- 103 Olsen KØ, Juul S, Bülow S, et al. Female fecundity before and after operation for familial adenomatous polyposis. *Br J Surg* 2003;90(02):227–231
- 104 Nelson H, Dozois RR, Kelly KA, Malkasian GD, Wolff BG, Ilstrup DM. The effect of pregnancy and delivery on the ileal pouch-anal anastomosis functions. *Dis Colon Rectum* 1989;32(05):384–388
- 105 Counihan TC, Roberts PL, Schoetz DJ Jr, Collier JA, Murray JJ, Veidenheimer MC. Fertility and sexual and gynecologic function after ileal pouch-anal anastomosis. *Dis Colon Rectum* 1994;37(11):1126–1129
- 106 Colwell JC, Gray M. What functional outcomes and complications should be taught to the patient with ulcerative colitis or familial adenomatous polyposis who undergoes ileal pouch anal anastomosis? *J Wound Ostomy Continence Nurs* 2001;28(04):184–189
- 107 Vrijland WW, Jeekel J, van Geldorp HJ, Swank DJ, Bonjer HJ. Abdominal adhesions: intestinal obstruction, pain, and infertility. *Surg Endosc* 2003;17(07):1017–1022
- 108 Milingos S, Kallipolitis G, Loutradis D, et al. Adhesions: laparoscopic surgery versus laparotomy. *Ann N Y Acad Sci* 2000;900:272–285
- 109 Asztély M, Palmblad S, Wikland M, Hultén L. Radiological study of changes in the pelvis in women following proctocolectomy. *Int J Colorectal Dis* 1991;6(02):103–107
- 110 Oresland T, Palmblad S, Ellström M, Berndtsson I, Crona N, Hultén L. Gynaecological and sexual function related to anatomical changes in the female pelvis after restorative proctocolectomy. *Int J Colorectal Dis* 1994;9(02):77–81
- 111 Juhasz ES, Fozard B, Dozois RR, Ilstrup DM, Nelson H. Ileal pouch-anal anastomosis function following childbirth. An extended evaluation. *Dis Colon Rectum* 1995;38(02):159–165
- 112 Metcalf AM, Dozois RR, Kelly KA. Sexual function in women after proctocolectomy. *Ann Surg* 1986;204(06):624–627
- 113 Nyam DC, Brilliant PT, Dozois RR, Kelly KA, Pemberton JH, Wolff BG. Ileal pouch-anal canal anastomosis for familial adenomatous polyposis: early and late results. *Ann Surg* 1997;226(04):514–519, discussion 519–521
- 114 Kartheuser AH, Parc R, Penna CP, et al. Ileal pouch-anal anastomosis as the first choice operation in patients with familial adenomatous polyposis: a ten-year experience. *Surgery* 1996;119(06):615–623